

l -ADIC MONODROMY AND SHIMURA CURVES IN POSITIVE CHARACTERISTICS

JIE XIA

ABSTRACT. In this paper, we seek an appropriate definition for a Shimura curve of Hodge type in positive characteristics, i.e. a characterization, in terms of geometry mod p , of curves in positive characteristics which are reduction of Shimura curve over $\mathbb{C}$. Specifically, we study the liftability of a curve in moduli space $\mathcal{A}_{4,1,k}$ of principally polarized abelian varieties over k , $\text{char } k = p$. We show that some conditions on the l -adic monodromy over such a curve imply that this curve can be lifted to a Shimura curve.

1. INTRODUCTION

This paper is a sequel to our previous paper [15] and [16]. All the three paper aim at answering the following question:

What is an appropriate definition of Shimura curves in positive characteristics ?

The answer is only known for Shimura varieties of PEL type which admit a natural moduli interpretation. In this paper, we consider Shimura curves of Hodge type and give an answer in the generic ordinary case, in terms of l -adic monodromy.

In [15], we start with a proper family of abelian varieties in characteristic p and prove if this family admits some special crystalline cycles, then it is a reduction of a Shimura curve of Mumford type. In this paper we present a similar result. We find the conditions on the l -adic monodromy associated to the family, which imply the family is a reduction of a Mumford type family of abelian fourfolds.

Let $\pi : X \rightarrow C$ be a family of principally polarized abelian fourfolds over a proper smooth curve C defined over a finite field $\mathbb{F}_q$ with $q = p^f$, $p > 3$. Let $\mathcal{E}_l$ be the lisse étale l -adic sheaf $R^1\pi_*(\mathbb{Q}_l)$ over C . Choose a geometric point $\bar{\xi}$ and a closed point c in C , and then $\mathcal{E}_l$ induces a monodromy:

$$\rho : \pi_1(C, \bar{\xi}) \rightarrow \text{Aut}(\mathcal{E}_{l,c}) \cong GL(8, \mathbb{Q}_l).$$

Let G_l be the Zariski closure of $\rho(\pi_1(C, \bar{\xi}))$ in $GL(8, \mathbb{Q}_l)$ and G_l^{geom} be that of $\rho(\pi_1^{\text{geom}}(C, \bar{\xi}))$.

$$1 \rightarrow \pi_1^{\text{geom}}(C, \bar{\xi}) \rightarrow \pi_1(C, \bar{\xi}) \rightarrow \text{Gal}(\bar{\mathbb{F}}_q/\mathbb{F}_q) \rightarrow 1.$$

Then G_l^{geom} is a normal subgroup of G_l .

To every closed point $c \in C$, it associates a unique (up to conjugation) Frobenius element F_c in $\pi_1(C, \bar{\xi})$. Its image in $\text{Gal}(\bar{\mathbb{F}}_q/\mathbb{F}_q)$ is the integer $\deg(\kappa(c) : \mathbb{F}_q)$.

For notational simplicity, let us define the following representation

$$\rho_0 : SL(2, \mathbb{Q}_l)^{\times 3} \rightarrow \text{Aut}(\mathcal{E}_{l,c})$$

as the tensor product of three copies of the standard representation of $SL(2, \mathbb{Q}_l)$.

Fix an embedding $\mathbb{Q}_l \rightarrow \mathbb{C}$ once for all. Our main theorem is as follows.

Theorem 1.1. *If there exists a closed point $c \in C$ such that*

- (1) $G_l^{\text{geom}, o} \otimes_{\mathbb{Q}_l} \mathbb{C} \cong \text{im } \rho_0 \otimes \mathbb{C}$,
- (2) X_c is an ordinary abelian variety,
- (3) In G_l , $\rho(F_c)$ generates a maximal torus which is unramified over $\mathbb{Q}_p$.

Then $X \rightarrow C$ is a weak Mumford curve.

If further assume the Higgs field of $X \rightarrow C$ is maximal, then there exists a family of polarized abelian fourfolds $Y \rightarrow C'$ such that

- (1) $C' \rightarrow C$ is a finite étale covering,
- (2) $Y \rightarrow X'$ is an isogeny between Y and the pullback family of X over C' ,
- (3) $Y \rightarrow C'$ is a good reduction of a Mumford curve.

For the definition of weak Mumford curve, see Section 2.

Remark 1.2. By 6.1 and Chebotarev density theorem, the Frobenius element over sufficiently many points in C generates the maximal torus in G_l . From [7], we know that the torus generated by the Frobenius F_c is defined over $\mathbb{Q}$. So it makes sense to require the torus is unramified over $\mathbb{Q}_p$, which is equivalent to say that the eigenvalues of the Frobenius are unramified over $\mathbb{Q}_p$.

We wonder if (3) can be replaced by a weaker condition, especially under the presence of (1) and (2). More are explained in Remark 6.2.

1.3. Structure of the paper. To prove 1.1, we reduce it to the main theorem in [15]. We need to compute the Frobenius eigenvalues in the l -adic setting and the connection with the crystalline cohomology.

In Section 3, we show some good reductions of Mumford curves satisfy the conditions in 1.1 so that we are not proving a vacuous theorem.

In Section 4, we compute the dimensions of the Frobenius eigenspaces in $H_{\text{et}}^0(C_{\mathbb{F}_q}, \wedge^4 \mathcal{E}_l)$ and $H_{\text{et}}^0(C_{\mathbb{F}_q}, \mathcal{E}nd(\wedge^2 \mathcal{E}_l))$.

$$(1.1) \quad \begin{aligned} \dim H_{\text{et}}^0(C_{\mathbb{F}_q}, \wedge^4 \mathcal{E}_l)^{F^{-q^2}} \otimes \mathbb{Q}_q &= 1 \\ \dim H_{\text{et}}^0(C_{\mathbb{F}_q}, \mathcal{E}nd(\wedge^2 \mathcal{E}_l))^F \otimes \mathbb{Q}_q &= 4. \end{aligned}$$

In Section 5, we translate the above result to p -adic case via comparing Lefschetz trace formulas. Therefore the Frobenius eigenspaces in crystalline cohomology spaces have the expected dimensions as in the main theorem of [15]. Furthermore, we conclude in Section 6 that

$$\Gamma((C/W(k))_{\text{cris}}, \mathcal{E}nd(\wedge^2 \mathcal{E}))^F \otimes \mathbb{Q}_q \cong \mathbb{Q}_q^{\times 4}$$

as algebras. Then 1.1 boils down to the main result in [15].

Acknowledgements. I am deeply indebted to my thesis advisor Aise Johan de Jong, who introduced me to this subject and encouraged me through this project. Without him I would have given up long time ago. I thank Martin Olsson for useful discussion and feedbacks.

2. MUMFORD CURVES

In [11, Chapter 4], Mumford defines a family of Shimura curves. We briefly recall the construction.

Let F be a cubic totally real field and D be a quaternion division algebra over F such that

$$D \otimes_{\mathbb{Q}} \mathbb{R} \cong \mathbb{H} \times \mathbb{H} \times M_2(\mathbb{R}), \text{Cor}_{F/\mathbb{Q}}(D) \cong M_8(\mathbb{Q}).$$

Here $\mathbb{H}$ is the quaternion algebra.

Let $G = \{x \in D^* | x\bar{x} = 1\}$. Then G is a $\mathbb{Q}$ -simple algebraic group and it is the $\mathbb{Q}$ -form of the $\mathbb{R}$ -algebraic group $SU(2) \times SU(2) \times SL(2, \mathbb{R})$.

$$h : \mathbb{S}_m(\mathbb{R}) \longrightarrow G(\mathbb{R})$$

$$e^{i\theta} \mapsto I_4 \otimes \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}.$$

The pair (G, h) forms a Shimura datum. And it defines Shimura curves, parameterizing abelian fourfolds. We call such curves (with its universal family) *Mumford curves*.

Definition 2.1. (1) A curve in $\mathcal{A}_{4,1,n} \otimes \mathbb{C}$ is called a *special Mumford curve* if it is the image of a Mumford curve in $\mathcal{A}_{4,1,n} \otimes \mathbb{C}$ induced by a universal family.
(2) The family $X \longrightarrow C$ is a *weak Mumford curve over k* if the image of $C \longrightarrow \mathcal{A}_{4,1,n}$ (induced by the family X/C) is (possible an irreducible component of) a reduction of a special Mumford curve in $\mathcal{A}_{4,1,n} \otimes \mathbb{C}$.

Remark 2.2. The “weakness” of the weak Mumford curve, comparing to good reductions, reflects in two aspects: firstly, the image of $C \longrightarrow \mathcal{A}_{4,1,n}$ might have singularities. Secondly, the reduction of a special Mumford curve at k might be reducible and that image of $C \longrightarrow \mathcal{A}_{4,1,n}$ is just one of the irreducible components.

3. EXAMPLES

To indicate that 1.1 is not a vacuous result, we show good reductions of a Mumford curve with an ordinary fiber satisfy the conditions of 1.1.

Let $f : A \longrightarrow M$ be the universal family over a Mumford curve defined over $\mathbb{C}$ and M is defined over the reflex field K . For every p over which K splits, $A \longrightarrow M$ admits a smooth and generically ordinary reduction $\pi : X \longrightarrow C$ over p ([9]). By the definition of Mumford curve or [14, Proposition 2.4], the image of

$$\rho_{\mathbb{C}} : \pi_1(M) \longrightarrow R^1 f_*(\underline{\mathbb{C}})$$

is $SL(2, \mathbb{C})^{\times 3}$. By Grothendieck specialization theorem of algebraic monodromy, $\rho_{\mathbb{C}}$ factors through

$$\rho : \pi_1(C, \bar{\xi}) \longrightarrow R^1 f_*(\underline{\mathbb{C}})$$

with a surjection $\pi_1(M) \longrightarrow \pi_1(\bar{C}, \bar{\xi})$. By the comparison of the l -adic cohomology and de Rham cohomology $R^1 f_*(\mathbb{C}) \cong R^1 f_*(\mathbb{Q}_l) \otimes \mathbb{C}$ for every $l \neq p$, we know conditions (1) and (2) in 1.1 hold for such C .

For condition (3), we choose x to be a CM point on M . Since A_x is simple, there is a degree 8 field $L \subset \text{End}^o(A_x)$. We can choose p such that L is unramified over p . Then look at the reduction $\bar{x}$ and $\mathbb{Q}[F_{\bar{x}}]$ is the center of $\text{End}(X_{\bar{x}})$. Since $L \subset \text{End}(X_{\bar{x}})$ is the maximal commutative subalgebra, $F \in L$ and in particular, F is unramified over p .

So with a careful choice of p , the resultant reduction of a Mumford curve satisfies all the conditions in 1.1.

4. FROBENIUS EIGENVALUES ON $\wedge^4 \mathcal{E}$ OR $\mathcal{E}nd(\wedge^2 \mathcal{E})$

In this section, we compute the dimension of Frobenius eigenspace at the target spaces and the corresponding eigenvalues.

Recall the short exact sequence

$$1 \longrightarrow \pi_1^{\text{geom}}(C, \bar{\xi}) \longrightarrow \pi_1(C, \bar{\xi}) \longrightarrow \text{Gal}(\bar{\mathbb{F}}_q/\mathbb{F}_q) \longrightarrow 1.$$

The Zariski closure G_l^{geom} , of $\rho(\pi_1^{\text{geom}}(C, \bar{\xi}))$ is a normal subgroup of G_l . Since $\mathcal{E}_l$ is pure, it follows from [4, 1.3.9, 3.4.1(iii)] that G_l^{geom} is semisimple. The connected component of identity $G_l^{\text{geom}, o}$ is the derived group of G_l^o .

Remark 4.1. If we assume $G_l^{\text{geom}} \otimes \mathbb{C}$ is entirely contained in $SL(2, \mathbb{C})^{\times 3}$, then we only need to enlarge the base field $\mathbb{F}_q$ to kill the S_3 part.

For every closed point c of C , the action of F_c on $\mathcal{E}_{l,c} \cong H_{\text{et}}^1(X_{\bar{c}}, \mathbb{Q}_l)$ is semisimple (see [2]). Therefore F_c acts on $\wedge^4 \mathcal{E}_{l,c}$ and $\text{End}(\wedge^2 \mathcal{E}_{l,c})$ semisimply.

Firstly let us consider the space

$$H_{\text{et}}^0(C_{\bar{\mathbb{F}}_q}, \mathcal{E}nd(\wedge^2 \mathcal{E}_l)) \cong \text{End}(\wedge^2(\mathcal{E}_{l,c}))^{\pi_1^{\text{geom}}(C, \bar{c})}.$$

Since F_c acts on $\text{End}(\wedge^2 \mathcal{E}_{l,c})$ semisimply, we can calculate its eigenvalues over $\mathbb{C}$.

Let V be the dimension 2 standard representation of $SL(2, \mathbb{C})$. Condition (1) of Theorem 1.1 shows that $\mathcal{E}_{l,c} \otimes \mathbb{C} \cong V^{\otimes 3}$ is the tensor product of three standard representations of $SL(2, \mathbb{C})$. So base change to $\mathbb{C}$,

$$H_{\text{et}}^0(C_{\bar{\mathbb{F}}_q}, \mathcal{E}nd(\wedge^2 \mathcal{E}_l)) \otimes \mathbb{C} \cong \text{End}(\wedge^2(V^{\otimes 3}))^{SL(2, \mathbb{C})^{\times 3}}.$$

As a representation of $SL(2, \mathbb{C})^{\times 3}$, $\wedge^2(V^{\otimes 3})$ decomposes into four distinct irreducible components

$$\wedge^2(V^{\otimes 3}) \cong \bigoplus_{i=1}^4 W_i.$$

There W_1, W_2, W_3 are all $S^2 V \otimes S^2 V$ and W_4 is the trivial representation of $SL(2, \mathbb{C})^{\times 3}$. Yet W_1, W_2, W_3 are pairwise non-isomorphic $SL(2, \mathbb{C})^{\times 3}$ representations.

Let p_k (i_k , resp.) be the projection from $\wedge^2(V^{\otimes 3})$ to W_k (inclusion from W_k to $\wedge^2(V^{\otimes 3})$, resp.). We have

$$\text{End}(\wedge^2(V^{\otimes 3}))^{SL(2, \mathbb{C})^{\times 3}} \cong \bigoplus_{k=1}^4 (i_k \circ \text{id}_{W_k} \circ p_k).$$

Note each id_{W_k} is invariant under the action of $SL(2, \mathbb{C})^{\times 3}$ and the scalar multiplication. Thus id_{W_k} is further invariant under the action of G_l . In particular, the Frobenius F_c fixes each id_{W_k} . So the invariant space $H_{\text{et}}^0(C_{\bar{\mathbb{F}}_q}, \mathcal{E}nd(\wedge^2 \mathcal{E}_l))^F$ has dimension 4.

Secondly,

$$H_{\text{et}}^0(C_{\bar{\mathbb{F}}_q}, \wedge^4 \mathcal{E}_l) = (\wedge^4 \mathcal{E}_{l,c})^{\pi_1^{\text{geom}}(C, \bar{\xi})}.$$

Similarly, base change to $\mathbb{C}$ and it is isomorphic to $\wedge^4(V^{\otimes 3})^{SL(2, \mathbb{C})^{\times 3}}$. One can directly compute by hand, or see the proof of Theorem 4.1 in [10] to conclude that this space only has dimension 1 which is generated by the polarization. Therefore the corresponding Frobenius eigenvalues are q^2 .

In summary, we have the following results:

$$(4.1) \quad \begin{aligned} \dim H_{\text{et}}^0(C_{\bar{\mathbb{F}}_q}, \mathcal{E}nd(\wedge^2 \mathcal{E}_l))^F &= 4, \\ \dim H_{\text{et}}^0(C_{\bar{\mathbb{F}}_q}, \wedge^4 \mathcal{E}_l)^{F-q^2} &= 1. \end{aligned}$$

5. COMPARISON OF LEFSCHETZ TRACE FORMULAS

In this section, we compare Lefschetz Trace Formulas to obtain a similar result to (4.1) in the case of crystalline cohomology.

We firstly consider $\mathcal{E}_p := R^1\pi_{\text{cris},*}(\mathcal{O}_X)$. Since σ is the identity on $\mathbb{F}_q$, the absolute Frobenius F acts linearly on $\mathcal{E}_{p,c}$. Since the local crystalline characteristic polynomial coincides with the l -adic one ([5, 1.3.5])

$$(5.1) \quad \det(1 - tF|_{\mathcal{E}_{p,c}}) = \det(1 - tF|_{\mathcal{E}_{l,c}}),$$

the eigenvalues of F on $\mathcal{E}_{l,c}$ and $\mathcal{E}_{p,c}$ are identical.

Let $\mathcal{F}_l$ be either $\wedge^4 \mathcal{E}_l$ or $\mathcal{E}nd(\wedge^2 \mathcal{E}_l)$. Since $\mathcal{E}_l$ comes from geometry, by Deligne's Weil II, the l -adic relative Lefschetz Trace Formula provides

$$\prod_{c \in C} \det(1 - tF|_{\mathcal{F}_{l,c}}) = \prod_i \det(1 - tF|_{H_{\text{et}}^i(C/\mathbb{F}_q, \mathcal{F}_l)})^{(-1)^i}.$$

In the p -adic case, we still use $\mathcal{F}_p$ to represent either $\wedge^4 \mathcal{E}_p$ or $\mathcal{E}nd(\wedge^2 \mathcal{E}_p)$. Since $\mathcal{E}_p$ is a Dieudonne crystal, $\mathcal{F}_p$ is automatically overconvergent. By a theorem of Etesse and le Stum ([6, 2.1.2]), we also have a Lefschetz Trace Formula within crystalline cohomology setting

$$\prod_{c \in C} \det(1 - tF|_{\mathcal{F}_{p,c}}) = \prod_i \det(1 - tF|_{H_{\text{cris}}^i(C/\mathbb{Z}_q, \mathcal{F}_p)})^{(-1)^i}.$$

Combining with equality (5.1), we have

$$(5.2) \quad \prod_i \det(1 - tF|_{H_{\text{et}}^i(C/\mathbb{F}_q, \mathcal{F}_l)})^{(-1)^i} = \prod_i \det(1 - tF|_{H_{\text{cris}}^i(C/\mathbb{Z}_q, \mathcal{F}_p)})^{(-1)^i}.$$

By Deligne's Weil II [4], the étale cohomology groups $H_{\text{et}}^i(C/\mathbb{F}_q, \mathcal{F}_l)$ is pure of weight $i + j$ where $\mathcal{F}_l$ has weight j . Since $\mathcal{F}_p$ is pointwisely pure, by [6, Theorem 5.3.2], $H_{\text{cris}}^i(C/\mathbb{Z}_q, \mathcal{F}_p)$ has the purity which implies, on each side of equality (5.2), there is no cancellation between the numerator and the denominator. All zeros or poles have the expected complex norms. Then we have the following termwise equality from (5.2).

$$(5.3) \quad \det(1 - tF|_{H_{\text{et}}^i(C/\mathbb{F}_q, \mathcal{F}_l)}) = \det(1 - tF|_{H_{\text{cris}}^i(C/\mathbb{Z}_q, \mathcal{F}_p)}).$$

So the eigenvalues of F on $H^0(C/\mathbb{Z}_q, \wedge^4 \mathcal{E}_p) \otimes \mathbb{Q}_q$ and $H^0(C/\mathbb{Z}_q, \mathcal{E}nd(\wedge^2 \mathcal{E}_p)) \otimes \mathbb{Q}_q$ are identical as on their l -adic counterparts. In particular,

$$(5.4) \quad \begin{aligned} \dim H^0(C/\mathbb{Z}_q, \wedge^4 \mathcal{E}_p)^{F-q^2} \otimes \mathbb{Q}_q &= 1, \\ \dim_{\mathbb{Q}_q} H^0(C/\mathbb{Z}_q, \mathcal{E}nd(\wedge^2 \mathcal{E}_p))^F \otimes \mathbb{Q}_q &= 4. \end{aligned}$$

6. COMPUTE $\text{End}^0(\wedge^2 \mathcal{E}_p)^F$

In order to apply the main theorem in [15], we need to prove that $\text{End}^0(\wedge^2 \mathcal{E}_p)^F \cong \mathbb{Q}_q^{\times 4}$ as algebras.

6.1. Frobenius Torus. By [13, Theorem 2], $\mathbb{Q}[F] \cong \prod K_i$ where K_i are number fields. The multiplicative group $\mathbb{Q}[F]^*$ defines a $\mathbb{Q}$ -torus

$$T = \prod_i \text{Res}_{K_i/\mathbb{Q}}(\mathbb{G}_m).$$

Viewing F as an element in G_l , T can be regarded as the $\mathbb{Q}$ -model of the connected component of 1 in the Zariski closure of the set $\{\rho(F)^n | n \in \mathbb{Z}\}$ in G_l (cf. [1, Chapter II, Section 13, Proposition 3]). In particular, T is contained in a maximal torus of G_l .

By [3, Theorem 3.7] and Chebotarev density theorem for the function field, generic points c on C satisfy that F_c generates a maximal torus. For every c , the torus T is defined over $\mathbb{Q}$. We say T is *unramified* over $\mathbb{Q}_p$ if the splitting field of T is unramified over prime p , and equivalently, the eigenvalues of F_c are unramified over p .

Remark 6.2. Varying the prime l , we obtain a compatible system of l -adic representation as stated in [7, 6.5]. The existence of a point c satisfying (3) in 1.1 requires that G_l is unramified over $\mathbb{Q}_p$.

On one hand, by [7, Proposition 8.9] and [8, Proposition 1.2, Theorem 3.2], for a subset of primes l of density 1 (or even l large enough), G_l is unramified over $\mathbb{Q}_l$. However, most results in the two paper have involved Dirichlet density restriction and hence can not be applied directly to our case.

On the other hand, we expect that if G_l is unramified over $\mathbb{Q}_q$, then there always exists a closed point c satisfying (3) in 1.1.

We also think generic ordinary property of $X \rightarrow C$ should also provide more information on Frobenius eigenvalues.

6.3. Eigenvalues of F_c on $\mathcal{E}_{pc}$. Note up to now, we have not used condition (2) and (3) in 1.1. Under the condition (2), by 6.1, we always can find c such that X_c ordinary and $\rho(F_c)$ a maximal torus. Further with the condition (3), there exists a closed point c which satisfies the following two conditions:

- (1) X_c is ordinary,
- (2) the Frobenius torus T is a maximal torus in G_l .

Now we study the eigenvalues of the Frobenius on the fiber over c . Since X_c is ordinary, $\mathcal{E}_{pc}$ is the product of a unit root crystal $\mathcal{U}_c$ and its dual $\mathcal{U}_c^\vee$. Let $\lambda_1, \dots, \lambda_4$ be the eigenvalues of F_c on $\mathcal{U}_c$. Then on $\mathcal{U}_c^\vee$, the eigenvalues of F_c are $\frac{q}{\lambda_i}$. Since $\mathcal{U}_c$ is a unit root crystal, λ_i are all p -adic units. Since $\mathcal{E}_{pc}$ has pure weight 1, λ_i all have complex norm $q^{\frac{1}{2}}$.

By 5.1, the Frobenius F_c also has eigenvalues $\lambda_1, \dots, \lambda_4, \frac{q}{\lambda_1}, \dots, \frac{q}{\lambda_4}$ on $\mathcal{E}_{l,c}$. Since the Frobenius torus is the maximal torus, the Frobenius eigenvalues λ_i correspond to the weights in the $SL(2)^{\times 3}$ representation $V^{\otimes 3}$. Let a, b, c be the three highest weights in the three standard representation of $SL(2, \mathbb{C})$. Then the eight weights of $V^{\otimes 3}$ are of the form $\pm a \pm b \pm c$ and they have a configuration as vertices of a cube. In this cube, the four p -adic units $\lambda_1, \dots, \lambda_4$ lie in the same face. Without loss of generality, we can assume that λ_1 corresponds to the highest weight $a + b + c$ and $\lambda_2, \lambda_3, \lambda_4$ correspond to $a + b - c, a + c - b$ and $a - b - c$. Then the only relation between $\lambda_1, \dots, \lambda_4$ is $\lambda_1 \lambda_4 = \lambda_2 \lambda_3$. So we have the following lemma.

Lemma 6.4. *Under the above choice of c , the eigenvalues λ_i have no relations other than those generated by $\lambda_i \frac{q}{\lambda_i} = q$ and $\lambda_1 \lambda_4 = \lambda_2 \lambda_3$.*

Remark 6.5. Lemma 6.4 also follows from the arguments in [12, Section 4].

Proposition 6.6.

$$\text{End}^0(\wedge^2 \mathcal{E}_p)^F \cong \mathbb{Q}_q^{\times 4}$$

as algebras.

Proof. From [15, Proposition 5.15] or basic representation theory of $SL(2)$, we know the condition 5.4 implies $\text{End}(\wedge^2 \mathcal{E}_p)^F \otimes_{\mathbb{Q}_q} \mathbb{C} \cong \mathbb{C}^{\times 4}$ as algebras. In particular, the algebra $\text{End}(\wedge^2 \mathcal{E}_p)^F$ is commutative. Therefore $\text{End}^0(\wedge^2 \mathcal{E}_p)^F$ is a product of fields.

Note $\wedge^2 \mathcal{E}_p$ has the polarization as a direct summand. So

$$\text{End}(\wedge^2 \mathcal{E}_p)^F \cong \mathbb{Q}_q^{\times 4}, \mathbb{Q}_q \times K \text{ or } \mathbb{Q}_q^{\times 2} \times L$$

where K is a degree 3 field extension of $\mathbb{Q}_q$ and L has degree 2. Comparing with the decomposition over $\mathbb{C}$, there exists $\eta_K \in K$ or $\eta_L \in L$ such that $\text{im } \eta_K$ and $\text{im } \eta_L$ are subcrystals in $\wedge^2 \mathcal{E}_p$. Further, $\text{rank im } \eta_K = 27$ and $\text{rank im } \eta_L = 18$.

If K or L is unramified over $\mathbb{Q}_q$, then by enlarging f in $q = p^f$, it becomes a product of copies of $\mathbb{Q}_q$. Therefore we only need to consider the case K or L ramified over $\mathbb{Q}_q$. Since $p \neq 2$ or 3 , we can assume $L \cong \mathbb{Q}_q(\sqrt{p})$ and $K \cong \mathbb{Q}_q(\sqrt[3]{p})$ and we can choose $\eta_K = \sqrt[3]{p}$, $\eta_L = \sqrt{p}$.

Note $\mathcal{E}_{pc} \cong \mathcal{U}_c \oplus \mathcal{U}_c^\vee$. Since the eigenvalues have distinct p -adic values, there is no F_c -invariant morphisms between $\wedge^2 \mathcal{U}_c$, $\wedge^2 \mathcal{U}_c^\vee$ and $\mathcal{U}_c \otimes \mathcal{U}_c^\vee$. Thus we have the decomposition

$$\text{End}(\wedge^2 \mathcal{E}_p)^F \longrightarrow \text{End}(\wedge^2 \mathcal{E}_{pc})^F \cong \text{End}(\wedge^2 \mathcal{U}_c) \oplus \text{End}(\wedge^2 \mathcal{U}_c^\vee) \oplus \text{End}(\mathcal{U}_c \otimes \mathcal{U}_c^\vee).$$

The restriction of F to $\text{End}(\wedge^2 \mathcal{E}_{pc})$ is just as F_c . Then by 6.4, all the eigenvalues of F on $\wedge^2 \mathcal{U}_c$ are $\lambda_1 \lambda_2, \dots, \lambda_3 \lambda_4$ and there is no more relations between the eigenvalues of $\wedge^2 \mathcal{U}_c$ other than $\lambda_4 \lambda_1 = \lambda_2 \lambda_3$. So each eigenspace $U_{\lambda_i \lambda_j}$ has dimension 1 except for $(1, 4)$ or $(2, 3)$. Thereby

$$(6.1) \quad \begin{aligned} \text{End}(\wedge^2 \mathcal{U}_c)^F &\cong \oplus_{(i,j) \neq (1,3), (2,4)} \text{End}(U_{\lambda_i \lambda_j}) \oplus \text{End}(U_{\lambda_1 \lambda_3}) \\ &\cong \oplus_{(i,j) \neq (1,3), (2,4)} \mathbb{Q}_q(\lambda_i \lambda_j) \oplus M_2(\mathbb{Q}_q(\lambda_1 \lambda_4)). \end{aligned}$$

Since the four eigenvalues λ_i are all unramified over $\mathbb{Q}_q$ and L or K is ramified, the image of the composition

$$L \text{ or } K \longrightarrow \text{End}(\wedge^2 \mathcal{E}_p)^F \longrightarrow \text{End}(\wedge^2 \mathcal{U}_c)^F$$

lies only in $\text{End}(U_{\lambda_1 \lambda_4}) \cong M_2(\mathbb{Q}_q(\lambda_1 \lambda_4))$. Otherwise, it would induce an embedding $L \text{ or } K \hookrightarrow \mathbb{Q}_q(\lambda_i \lambda_j)$. In particular, $\eta_K|_{\wedge^2 \mathcal{U}_c}$ or $\eta_L|_{\wedge^2 \mathcal{U}_c}$ has only rank 2.

Restricted to point c , the image of η_K has dimension at most only 20. Contradiction.

For L , we know that $\eta_L|_{\mathcal{U}_c \otimes \mathcal{U}_c^\vee}$ is a surjection. Note the eigenvalues of F_c on $\mathcal{U}_c \otimes \mathcal{U}_c^\vee$ have the form $\frac{q\lambda_i}{\lambda_j}$. Again by 6.4, among these eigenvalues, $\frac{q\lambda_1}{\lambda_4}$ has only multiplicity 1.

Therefore

$$\text{End}(\mathcal{U}_c \otimes \mathcal{U}_c^\vee)^F \cong \text{End}(\mathcal{U}_{\frac{q\lambda_1}{\lambda_4}}) \oplus \dots \cong \mathbb{Q}_q(\frac{q\lambda_1}{\lambda_4}) \oplus \dots$$

as algebras. Since $\mathbb{Q}_q(\frac{q\lambda_1}{\lambda_4})$ is unramified over $\mathbb{Q}_q$, the image of L in $\text{End}(\mathcal{U}_c \otimes \mathcal{U}_c^\vee)^F$ excludes $\text{End}(\mathcal{U}_{\frac{q\lambda_1}{\lambda_4}})$ and hence η_L can not be a surjection. The contradiction concludes the proof. $\square$

REFERENCES

1. Claude Chevalley, *Théorie des groupes de Lie. Tome II. Groupes algébriques*, Actualités Sci. Ind. no. 1152, Hermann & Cie., Paris, 1951. MR 0051242 (14,448d)
2. Wên Chên Chi, *On the l -adic representations attached to some absolutely simple abelian varieties of type II*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **37** (1990), no. 2, 467–484. MR 1071431 (91g:11060)
3. ———, *l -adic and λ -adic representations associated to abelian varieties defined over number fields*, Amer. J. Math. **114** (1992), no. 2, 315–353. MR 1156568 (93c:11038)
4. Pierre Deligne, *La conjecture de Weil. II*, Inst. Hautes Études Sci. Publ. Math. (1980), no. 52, 137–252. MR 601520 (83c:14017)
5. Luc Illusie, *Crystalline cohomology*, Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., vol. 55, Amer. Math. Soc., Providence, RI, 1994, pp. 43–70. MR 1265522 (95a:14021)
6. Kiran S. Kedlaya, *Fourier transforms and p -adic ‘Weil II’*, Compos. Math. **142** (2006), no. 6, 1426–1450. MR 2278753 (2008b:14024)
7. M. Larsen and R. Pink, *On l -independence of algebraic monodromy groups in compatible systems of representations*, Invent. Math. **107** (1992), no. 3, 603–636. MR 1150604 (93h:22031)
8. ———, *Abelian varieties, l -adic representations, and l -independence*, Math. Ann. **302** (1995), no. 3, 561–579. MR 1339927 (97e:14057)
9. Sheng Mao and Kang Zuo, *On the newton polygons of abelian varieties of mumford’s type*, arXiv:1106.3505v1.
10. B. J. J. Moonen and Yu. G. Zarhin, *Hodge classes and Tate classes on simple abelian fourfolds*, Duke Math. J. **77** (1995), no. 3, 553–581. MR 1324634 (96b:14010)
11. D. Mumford, *A note of Shimura’s paper “Discontinuous groups and abelian varieties”*, Math. Ann. **181** (1969), 345–351. MR 0248146 (40 #1400)
12. Rutger Noot, *Abelian varieties with l -adic Galois representation of Mumford’s type*, J. Reine Angew. Math. **519** (2000), 155–169. MR 1739726 (2001k:11112)
13. John Tate, *Endomorphisms of abelian varieties over finite fields*, Invent. Math. **2** (1966), 134–144. MR 0206004 (34 #5829)
14. Jie Xia, *A characterization of the good reduction of Mumford curve*, arXiv: 1306.0163.
15. Jie Xia, *Crystalline Hodge cycles and Shimura curves in positive characteristic*, arXiv:1311.0940.
16. Jie Xia, *Tensor decomposition of isocrystals characterizes Mumford curves*, arXiv:1310.2682.

MATHEMATICS DEPARTMENT, COLUMBIA UNIVERSITY IN THE CITY OF NEW YORK
E-mail address: xiajie@math.columbia.edu